



Social Distancing & Self Isolation Guidelines

Updated 17th March 2020

Source- <https://www.gov.uk/government/publications/covid-19-guidance-on-social-distancing-and-for-vulnerable-people/guidance-on-social-distancing-for-everyone-in-the-uk-and-protecting-older-people-and-vulnerable-adults>

The UK Govt are advising those who are at increased risk of severe illness from coronavirus (COVID-19) to be particularly stringent in following social distancing measures.

This group includes those who are:

- aged 70 or older (regardless of medical conditions)
- under 70 with an underlying health condition listed below (ie anyone instructed to get a flu jab as an adult each year on medical grounds):
 - chronic (long-term) respiratory diseases, such as [asthma](#), [chronic obstructive pulmonary disease \(COPD\)](#), emphysema or [bronchitis](#)
 - chronic heart disease, such as [heart failure](#)
 - [chronic kidney disease](#)
 - chronic liver disease, such as [hepatitis](#)
 - chronic neurological conditions, such as [Parkinson's disease](#), [motor neurone disease](#), [multiple sclerosis \(MS\)](#), a learning disability or cerebral palsy
 - [diabetes](#)
 - problems with your spleen – for example, [sickle cell](#) disease or if you have had your spleen removed
 - a weakened immune system as the result of conditions such as [HIV and AIDS](#), or medicines such as [steroid tablets](#) or [chemotherapy](#)
 - being seriously overweight (a body mass index (BMI) of 40 or above)
- those who are pregnant

Note: there are some clinical conditions which put people at even higher risk of severe illness from COVID-19. If you are in this category, next week the NHS in England will directly contact you with advice the more stringent measures you should take in order to keep yourself and others safe. For now, you should rigorously follow the social distancing advice in full, outlined below.

People falling into this group are those who may be at particular risk due to complex health problems such as:

- people who have received an organ transplant and remain on ongoing immunosuppression medication
- people with cancer who are undergoing active chemotherapy or radiotherapy
- people with cancers of the blood or bone marrow such as leukaemia who are at any stage of treatment
- people with severe chest conditions such as cystic fibrosis or severe asthma (requiring hospital admissions or courses of steroid tablets)
- people with severe diseases of body systems, such as severe kidney disease (dialysis)

What is social distancing?

Social distancing measures are steps you can take to reduce the social interaction between people. This will help reduce the transmission of coronavirus (COVID-19).

They are:

1. Avoid contact with someone who is displaying symptoms of coronavirus (COVID-19). These symptoms include high temperature and/or new and continuous cough
2. Avoid non-essential use of public transport, varying your travel times to avoid rush hour, when possible
3. Work from home, where possible. Your employer should support you to do this. Please refer to [employer guidance](#) for more information
4. Avoid large gatherings, and gatherings in smaller public spaces such as pubs, cinemas, restaurants, theatres, bars, clubs
5. Avoid gatherings with friends and family. Keep in touch using remote technology such as phone, internet, and social media
6. Use telephone or online services to contact your GP or other essential services

Everyone should be trying to follow these measures as much as is pragmatic.

For those who are over 70, have an underlying health condition or are pregnant, we strongly advise you to follow the above measures as much as you can, and to significantly limit your face-to-face interaction with friends and family if possible.

This advice is likely to be in place for some weeks.

Handwashing and respiratory hygiene

There are general principles you can follow to help prevent the spread of respiratory viruses, including:

- washing your hands more often - with soap and water for at least 20 seconds or use a hand sanitiser when you get home or into work, when you blow your nose, sneeze or cough, eat or handle food

- avoid touching your eyes, nose, and mouth with unwashed hands
- avoid close contact with people who have symptoms
- cover your cough or sneeze with a tissue, then throw the tissue in a bin and wash your hands
- clean and disinfect frequently touched objects and surfaces in the home

Please see – <https://www.gov.uk/government/publications/covid-19-guidance-on-social-distancing-and-for-vulnerable-people/guidance-on-social-distancing-for-everyone-in-the-uk-and-protecting-older-people-and-vulnerable-adults> for further information

Self Isolation Guidance –

Updated 17th March 2020

Source – <https://www.acas.org.uk/coronavirus>

Self-isolation and sick pay

Employees and workers must receive any [Statutory Sick Pay \(SSP\)](#) due to them if they need to self-isolate because:

- they have coronavirus
- they have coronavirus symptoms, for example a high temperature or new continuous cough
- someone in their household has coronavirus symptoms
- they've been told to self-isolate by a doctor or NHS 111

If someone has symptoms, everyone in their household must self-isolate for 14 days.

Employers might offer more than SSP – 'contractual' sick pay. [Find out more about sick pay.](#)

If an employee or worker cannot work, they should tell their employer:

- as soon as possible
- the reason
- how long they're likely to be off for

The employer might need to be flexible if they require evidence from the employee or worker. For example, someone might not be able to provide a [sick note \('fit note'\)](#) if they've been told to self-isolate for more than 7 days.

For more information please go to - <https://www.acas.org.uk/coronavirus>